


Just to be clear about Free Energy



## Last Phase change

What is a key difference between evaporation and boiling?

- A. liquids only boil at 1 atm total pressure
- B. liquids only evaporate at room temperature
- C. bubble form in liquids when boiling
- D. nothing

## Boiling demo

## Solutions


Solutions are homogeneous mixtures of multiple compounds

Solution  
salt water  
air  
steel

Major component = Solvent  
(language typically used for liquids)

Minor component = Solute


Let's look at the following "reaction"



Which has the higher entropy?

- A. The water + the solid salt
- B. The solution ←
- C. They are exactly the same

Let's look at the following "reaction"



Which has the higher enthalpy?

- A. The water + the solid salt
- B. The solution
- C. They are essentially the same ←

Let's look at the following "reaction"



Which has the lower free energy?

- A. The water + the solid salt
- B. The solution ←
- C. They are exactly the same

What is enthalpy change for making a solution?

Lose solute-solute interactions (IMF)  
Lose solvent-solvent interactions (IMF)  
Gain solute-solvent interactions




Figure Copyright Houghton Mifflin Company. All rights reserved

## Enthalpy of Solvation $\Delta H_{\text{solvation}}$ hard to predict

$$\Delta H_{\text{solvation}} = 0$$

Ideal solution

Solute-solvent interactions are identical to  
solute-solute and solvent-solvent

$$\Delta H_{\text{solvation}} > 0$$

Typical

Solute-solvent interactions are weaker than  
solute-solute and solvent-solvent

$$\Delta H_{\text{solvation}} < 0$$

Unusual but possible

Solute-solvent interactions are stronger than  
solute-solute and solvent-solvent

## Solvation Demo

For the dissolution of ammonium nitrate  
the free energy

- A. decreases
- B. increases
- C. stays the same



## Entropy of Solvation $\Delta S_{\text{solvation}}$ usually easy to predict

Solutions have a higher entropy than the unmixed compounds

Therefore

$$\Delta S_{\text{solvation}} > 0$$

For most cases

Not true for small high charge density ions

**TABLE 17.2** Values of  $\Delta S_{\text{soln}}^{\circ}$  for Several Salts Dissolving in Water

Process	$\Delta S^{\circ}$ (J K <sup>-1</sup> mol <sup>-1</sup> )
KCl(s) → K <sup>+</sup> (aq) + Cl <sup>-</sup> (aq)	75
LiF(s) → Li <sup>+</sup> (aq) + F <sup>-</sup> (aq)	-36
CaS(s) → Ca <sup>2+</sup> (aq) + S <sup>2-</sup> (aq)	-138

### Gibb's Free Energy of Solvation $\Delta G_{\text{solvation}}$

If  $\Delta G_{\text{solvation}} < 0$  solution strongly favored

If  $\Delta G_{\text{solvation}} > 0$  undissolved state is strongly favored

$$\Delta G_{\text{solvation}} = \Delta H_{\text{solvation}} - T \Delta S_{\text{solvation}}$$

Typically  $\Delta S_{\text{solvation}} > 0$ ,  $\Delta H_{\text{solvation}} > 0$

need  $|T\Delta S| > |\Delta H|$

What makes an ideal solution?

Same IMF for solute-solvent and solute-solute and solvent-solvent

**"like dissolves like"**

Polar compounds dissolve polar compounds (ionic)

Nonpolar compound dissolve nonpolar compounds

This minimize  $\Delta H$

### Miscibility Demo

Definitions:

Miscible: capable of being mixed  
Immiscible: incapable of being mixed

Which is most likely to dissolve best in water?


- A. methanol  $\text{CH}_3\text{OH}$  ←
- B. butanol  $\text{C}_4\text{H}_9\text{OH}$
- C. octanol  $\text{C}_8\text{H}_{17}\text{OH}$
- D. didodecanol  $\text{C}_{12}\text{H}_{25}\text{OH}$

Which is most likely to dissolve best in hexane ( $\text{C}_6\text{H}_{14}$ )?

- A. methanol  $\text{CH}_3\text{OH}$
- B. butanol  $\text{C}_4\text{H}_9\text{OH}$
- C. octanol  $\text{C}_8\text{H}_{17}\text{OH}$
- D. didodecanol  $\text{C}_{12}\text{H}_{25}\text{OH}$  ←

### Temperature Dependence

Generally at T goes up solubility increases



### Gas Dissolved in a Liquid

Henry's Law

**TABLE 17.3** The Values of Henry's Law Constants for Several Gases Dissolved in Water at 298 K

Gas	$k_H$ (atm)
$\text{CH}_4$	$4.13 \times 10^2$
$\text{CO}_2$	$1.64 \times 10^3$
$\text{O}_2$	$4.34 \times 10^4$
$\text{CO}$	$5.71 \times 10^4$
$\text{H}_2$	$7.03 \times 10^4$
$\text{N}_2$	$8.57 \times 10^4$

$$P_{\text{solute}} = K_{\text{solvent}} X_{\text{solute}}$$

↑  
mole fraction

## In General

Henry's Law constants increase with increasing Temperature

Less gas is dissolved at higher temperatures

$$\Delta H < 0$$

going from no attractions to being in a liquid