Today


Quantitative Echem How many electrons

Qualitative Echem
Which direction is spontaneous
What is the voltage
What is the best "oxidizer"

First some nomenclature

Galvanic Cell Voltaic Cell Battery

Spontaneous $\Delta G < 0$ E > 0



Cathode at a higher potential than the anode So cathode get the PLUS sign


This is spontaneous. It can be used as a power supply

Principles of Chemistry II

First some nomenclature

Electrolytic Cell

Non-Spontaneous $\Delta G > 0$ E < 0



Anode at a higher potential than the cathode So anode get the PLUS sign


This reaction must be driven by an external power supply

In the following standard Ecell, what is the sign of the cathode?

$$Zn(s) | Zn^{2+} | | H^{+} | H_{2} (g)$$

 $Zn^{2+} + 2e^{-} ---> Zn(s) E^{\circ} = -0.76 V$
 $2H^{+} + 2e^{-} ---> H_{2} E^{\circ} = 0.0 V$

- A +
- B. -
- C. neither E° cell = 0



Oxidation Cathode Anode Anode Zn²⁺ SO₄²⁻ Zn(s) --Cu(s) $.0 M \text{ Cu}^{2^{-1}}$ solution Zn 2+

Reduction Cathode

We've made a I.I V battery!

If I use this battery for a while how much Zn reacts?

Coulomb (C) = Amp (C
$$s^{-1}$$
) x Second (s)

How many electrons are in a Coulomb? What is the charge of I mole of electrons?

F is the charge of one mole of electrons $F = 96,485 \, \text{C}$ (Faraday's Constant)

If I run this cell for 100 s at a current of 30 mA how many moles of electrons flow?

A.
$$(30 \times 10^{-3}) \times 100 \times F$$

$$Q = A \times t = 10 \times 10^{-3} A \times 100 s$$

B.
$$30 \times 100 \times F$$

C.
$$30 / (100 \times F)$$

D.
$$(30 \times 10^{-3}) / (100 \times F)$$

E.
$$[(30 \times 10^{-3}) \times 100] / F$$

$$= 3 / F$$

If I run this cell for 100 s at a current of I mA how many moles of Zn react?

$$Zn(s) | Zn^{2+} || Cu^{2+} | Cu (s)$$

A. (3/F)

B. $(3/F) \times 2$

C. (3/F) / 2


I mole of Zn²⁺ requires
2 electron to be
reduced to Zn

So 3/F moles of electrons will produce half that number of moles of Zn

$$Zn(s) + Cu^{2+}(aq) \longrightarrow Zn^{2+}(aq) + Cu(s)$$

 $Zn(s) \mid Zn^{2+} \mid \mid Cu^{2+} \mid Cu$


Oxidation Anode




Reduction Cathode

We've made a I.I V battery!

How do we know what the voltage is?



The voltage depends on the concentrations (we've all had dead batteries)



Mix up "standard" concentrations
I M Zn²⁺ and I M Cu²⁺
(note this is very concentrated)

Now we can measure every possible combination of electrochemical cells!

What if I would like to predict the voltage from a cell using the following reaction at standard conditions?

$$Mn^{2+} + I_2 \longrightarrow MnO_4^- + I^-$$

If want to know about the potential we just have to identify the two reactions!

$$Mn^{2+} + I_2 \longrightarrow MnO_4^- + I^-$$

$$|_2 \longrightarrow |_{-}$$

$$|_2 \longrightarrow 2|$$

$$I_2 + 2e^- \longrightarrow 2I^-$$
 This is the reduction (Cathode)

$$Mn^{2+} + I_2 \longrightarrow MnO_4^- + I^-$$

$$Mn^{2+} \longrightarrow MnO_4^-$$

everything but O & H

$$4H_2O + Mn^{2+} \longrightarrow MnO_4^-$$

balance O w/ H₂O

$$4H_2O + Mn^{2+} \longrightarrow MnO_4^- + 8H^+$$
 balance H w/ H⁺

$$4H_2O + Mn^{2+} \longrightarrow MnO_4^- + 8H^+ + 5 e^-$$
 charge w/e

This is the oxidation (Anode)

$$Mn^{2+} + I_2 \longrightarrow MnO_4^- + I^-$$

$$4H_2O + Mn^{2+} \longrightarrow MnO_4^- + 8H^+ + 5 e^-$$

 $I_2 + 2e^- \longrightarrow 2I^-$

To compare these two we don't need to balance the number of electrons!

We only need to do this to relate moles of electrons to moles of materials.


Potential difference depends only on the two reactions!


$$4H_2O + Mn^{2+} \longrightarrow MnO_4^- + 8H^+ + 5 e^-$$

 $I_2 + 2e^- \longrightarrow 2I^-$

How can we compare these two?

We'll compare every reaction to a standard






But if a hole appears beneath me then it is no longer zero and I will move spontaneously from high to low!

We need to pick a zero potential for electrochemistry

We chose this reaction



$$2H^+ + 2e^- H_2(g)$$


note standard conditions

we pick this as E°= 0V

potential energy

Now compare everything to this

Principles of Chemistry II



So potential for
$$Zn \longrightarrow Zn^{2+} + 2e^{-}$$
 is 0.76 V

Write everything as a reduction reaction

Half-reaction	\mathscr{E}° (V)	Half-reaction	&° (V)
$F_2 + 2e^- \rightarrow 2F^-$	2.87	$O_2 + 2H_2O + 4e^- \rightarrow 4OH^-$	0.40
$Ag^{2+} + e^- \rightarrow Ag^+$	1.99	$Cu^{2+} + 2e^{-} \rightarrow Cu$	0.34
$Co^{3+} + e^- \rightarrow Co^{2+}$	1.82	$Hg_2Cl_2 + 2e^- \rightarrow 2Hg + 2Cl^-$	0.27
$H_2O_2 + 2H^+ + 2e^- \rightarrow 2H_2O$	1.78	$AgCl + e^- \rightarrow Ag + Cl^-$	0.22
$Ce^{4+} + e^{-} \rightarrow Ce^{3+}$	1.70	$SO_4^{2-} + 4H^+ + 2e^- \rightarrow H_2SO_3 + H_2O$	0.20
$PbO_2 + 4H^+ + SO_4^{2-} + 2e^- \rightarrow PbSO_4 + 2H_2O$	1.69	$Cu^{2+} + e^{-} \rightarrow Cu^{+}$	0.16
$MnO_4^- + 4H^+ + 3e^- \rightarrow MnO_2 + 2H_2O$	1.68	$2H^+ + 2e^- \rightarrow H_2$	0.00
$IO_4^- + 2H^+ + 2e^- \rightarrow IO_3^- + H_2O$	1.60	$Fe^{3+} + 3e^{-} \rightarrow Fe$	-0.03
$MnO_4^- + 8H^+ + 5e^- \rightarrow Mn^{2+} + 4H_2O$	1.51	$Pb^{2+} + 2e^{-} \rightarrow Pb$	-0.13
$Au^{3+} + 3e^- \rightarrow Au$	1.50	$\mathrm{Sn}^{2+} + 2\mathrm{e}^{-} \rightarrow \mathrm{Sn}$	-0.14
$PbO_2 + 4H^+ + 2e^- \rightarrow Pb^{2+} + 2H_2O$	1.46	$Ni^{2+} + 2e^- \rightarrow Ni$	-0.23
$Cl_2 + 2e^- \rightarrow 2Cl^-$	1.36	$PbSO_4 + 2e^- \rightarrow Pb + SO_4^{2-}$	-0.35
$Cr_2O_7^{2-} + 14H^+ + 6e^- \rightarrow 2Cr^{3+} + 7H_2O$	1.33	$Cd^{2+} + 2e^{-} \rightarrow Cd$	-0.40
$O_2 + 4H^+ + 4e^- \rightarrow 2H_2O$	1.23	$Fe^{2+} + 2e^{-} \rightarrow Fe$	-0.44
$MnO_2 + 4H^+ + 2e^- \rightarrow Mn^{2+} + 2H_2O$	1.21	$Cr^{3+} + e^- \rightarrow Cr^{2+}$	-0.50
$IO_3^- + 6H^+ + 5e^- \rightarrow \frac{1}{2}I_2 + 3H_2O$	1.20	$Cr^{3+} + 3e^- \rightarrow Cr$	-0.73
$Br_2 + 2e^- \rightarrow 2Br^-$	1.09	$Zn^{2+} + 2e^- \rightarrow Zn$	-0.76
$VO_2^+ + 2H^+ + e^- \rightarrow VO^{2+} + H_2O$	1.00	$2H_2O + 2e^- \rightarrow H_2 + 2OH^-$	-0.83
$AuCl_4^- + 3e^- \rightarrow Au + 4Cl^-$	0.99	$Mn^{2+} + 2e^- \rightarrow Mn$	-1.18
$NO_3^- + 4H^+ + 3e^- \rightarrow NO + 2H_2O$	0.96	$Al^{3+} + 3e^- \rightarrow Al$	-1.66
$ClO_2 + e^- \rightarrow ClO_2^-$	0.954	$H_2 + 2e^- \rightarrow 2H^-$	-2.23
$2Hg^{2+} + 2e^{-} \rightarrow Hg_{2}^{2+}$	0.91	$Mg^{2+} + 2e^- \rightarrow Mg$	-2.37
$Ag^+ + e^- \rightarrow Ag$	0.80	$La^{3+} + 3e^- \rightarrow La$	-2.37
$Hg_2^{2+} + 2e^- \rightarrow 2Hg$	0.80	$Na^+ + e^- \rightarrow Na$	-2.71
$Fe^{3+} + e^- \rightarrow Fe^{2+}$	0.77	$Ca^{2+} + 2e^{-} \rightarrow Ca$	-2.76
$O_2 + 2H^+ + 2e^- \rightarrow H_2O_2$	0.68	$Ba^{2+} + 2e^{-} \rightarrow Ba$	-2.90
$MnO_4^- + e^- \rightarrow MnO_4^{2-}$	0.56	$K^+ + e^- \rightarrow K$	-2.92
$I_2 + 2e^- \rightarrow 2I^-$ $Cu^+ + e^- \rightarrow Cu$	0.54	$Li^+ + e^- \rightarrow Li$	-3.05

$\begin{array}{lll} & \text{Easy to} \\ +2.87 & F_2(s) + 2e^- \longrightarrow 2F^-(sq) \\ +1.51 & \text{MnO}_4^-(sq) + 8\text{H}^+(sq) + 5e^- \longrightarrow \text{Mn}^2+(sq) + 4\text{H}_2\text{O}(/) \\ +1.36 & \text{Cl}_2(s) + 2e^- \longrightarrow 2\text{Cl}^-(sq) \\ +1.33 & \text{Cr}_2\text{O}_7^{2-}(sq) + 14\text{H}^+(sq) + 6e^- \longrightarrow 2\text{Cr}^{3+}(sq) + 7\text{H}_2\text{O}(/) \\ +1.23 & \text{O}_2(s) + 4\text{H}^+(sq) + 4e^- \longrightarrow 2\text{H}_2\text{O}(/) \\ +1.06 & \text{Br}_2(/) + 2e^- \longrightarrow 2\text{Br}^-(sq) \\ +0.96 & \text{NO}_3^-(sq) + 4\text{H}^+(sq) + 3e^- \longrightarrow \text{NO}(s) + \text{H}_2\text{O}(/) \\ +0.80 & \text{A}_8^+(sq) + e^- \longrightarrow \text{A}_8(s) \\ +0.77 & \text{Fe}^{3+}(sq) + e^- \longrightarrow \text{Fe}^{2+}(sq) \\ \end{array}$	ngest izing
+1.51 $MnO_4^-(sq) + 8H^+(sq) + 5e^- \longrightarrow Mn^2 + (sq) + 4H_2O(I)$ +1.36 $Cl_2(g) + 2e^- \longrightarrow 2Cl^-(sq)$ +1.33 $Cr_2O_7^{2-}(sq) + 14H^+(sq) + 6e^- \longrightarrow 2Cr^{3+}(sq) + 7H_2O(I)$ +1.23 $O_2(g) + 4H^+(sq) + 4e^- \longrightarrow 2H_2O(I)$ +1.06 $Br_2(I) + 2e^- \longrightarrow 2Br^-(sq)$ +0.96 $NO_3^-(sq) + 4H^+(sq) + 3e^- \longrightarrow NO(g) + H_2O(I)$ +0.80 $Ag^+(sq) + e^- \longrightarrow Ag(s)$ +0.77 $Fe^{3+}(sq) + e^- \longrightarrow Fe^{2+}(sq)$	izing
+1.36 $Cl_2(g) + 2e^- \longrightarrow 2Cl^-(sq)$ $Cl_2(g) + 2e^- \longrightarrow 2Cl^-(sq)$ $Cr_2O_7^{2-}(sq) + 14H^+(sq) + 6e^- \longrightarrow 2Cr^{3+}(sq) + 7H_2O(I)$ +1.23 $O_2(g) + 4H^+(sq) + 4e^- \longrightarrow 2H_2O(I)$ ager +1.06 $Br_2(I) + 2e^- \longrightarrow 2Br^-(sq)$ $Pl_2(G) + Pl_2(G) + Pl$	izing
+1.33 $Cr_2O_7^{2-}(sq) + 14H^+(sq) + 6e^- \longrightarrow 2Cr^{3+}(sq) + 7H_2O(I)$ +1.23 $O_2(g) + 4H^+(sq) + 4e^- \longrightarrow 2H_2O(I)$ ager +1.06 $Br_2(I) + 2e^- \longrightarrow 2Br^-(sq)$ +0.96 $NO_3^-(sq) + 4H^+(sq) + 3e^- \longrightarrow NO(g) + H_2O(I)$ +0.80 $Ag^+(sq) + e^- \longrightarrow Ag(s)$ +0.77 $Fe^{3+}(sq) + e^- \longrightarrow Fe^{2+}(sq)$	
+1.33 $Cr_2O_7^{2-}(sq) + 14H^+(sq) + 6e^- \longrightarrow 2Cr^{3+}(sq) + 7H_2O(I)$ +1.23 $O_2(g) + 4H^+(sq) + 4e^- \longrightarrow 2H_2O(I)$ ager +1.06 $Br_2(I) + 2e^- \longrightarrow 2Br^-(sq)$ +0.96 $NO_3^-(sq) + 4H^+(sq) + 3e^- \longrightarrow NO(g) + H_2O(I)$ +0.80 $Ag^+(sq) + e^- \longrightarrow Ag(s)$ +0.77 $Fe^{3+}(sq) + e^- \longrightarrow Fe^{2+}(sq)$	
+1.06 $Br_2(I) + 2e^- \longrightarrow 2Br^-(sq)$ +0.96 $NO_3^-(sq) + 4H^+(sq) + 3e^- \longrightarrow NO(g) + H_2O(I)$ +0.80 $Ag^+(sq) + e^- \longrightarrow Ag(s)$ +0.77 $Fe^{3+}(sq) + e^- \longrightarrow Fe^{2+}(sq)$	TTS)
+0.96 $NO_3^-(sq) + 4H^+(sq) + 3e^- \longrightarrow NO(g) + H_2O(f)$ +0.80 $Ag^+(sq) + e^- \longrightarrow Ag(s)$ +0.77 $Fe^{3+}(sq) + e^- \longrightarrow Fe^{2+}(sq)$,
+0.80 $Ag^+(sq) + e^- \longrightarrow Ag(s)$ +0.77 $Fe^{3+}(sq) + e^- \longrightarrow Fe^{2+}(sq)$	
$+0.77$ $Fe^{3+}(sq) + e^{-} \longrightarrow Fe^{2+}(sq)$	
$+0.68$ $O_2(g) + 2H^+(aq) + 2e^- \longrightarrow H_2O_2(aq)$	
$+0.59$ $MnO_4^-(sq) + 2H_2O(I) + 3e^- \longrightarrow MnO_2(s) + 4OH^-(sq)$	
$+0.54$ $I_2(s) + 2e^- \longrightarrow 2I^-(sq)$	
$+0.40$ $O_2(g) + 2H_2O(I) + 4e^- \longrightarrow 4OH^-(sq)$	
$+0.34$ $Cu^{2+}(sq) + 2e^{-} \longrightarrow Cu(s)$	
$0 2H^{+}(sq) + 2e^{-} \longrightarrow H_{2}(s)$	
-0.28 $Ni^{2+}(sq) + 2e^{-} \longrightarrow Ni(s)$	
-0.44 Fe ²⁺ (sq) + 2e ⁻ \longrightarrow Fe(s)	oxidize
-0.76 $Zrr^{2+}(sq) + 2e^{-} \longrightarrow Zr(s)$ (ctror	agost
$\begin{array}{ccc} -0.76 & & ZH^{-1}(sq) + Ze^{-1} \longrightarrow ZH(s) \\ -0.83 & & 2H_2O(s) + 2e^{-1} \longrightarrow H_2(s) + 2OH^{-1}(sq) \end{array} $ (strong)	igest
-1.66 $Al^{3+}(sq) + 3e^{-} \longrightarrow Al(s)$ redu	cing
-2.71 Na ⁺ (sq) + e ⁻ \longrightarrow Ns(s)	, o
-3.05 $\text{Li}^{+}(sq) + e^{-} \longrightarrow \text{Li}(s)$ ager	its)

How to find E°cell?

$$E^{\circ}_{cell} = E^{\circ}_{cathode} - E^{\circ}_{anode}$$

$$4H_2O + Mn^{2+} \longrightarrow MnO_4^- + 8H^+ + 5 e^ I_2 + 2e^- \longrightarrow 2I^ I_2 + 2e^- \longrightarrow 2I^ E^\circ = 0.54V$$
 $MnO_4^- + 8H^+ + 5 e^- \longrightarrow 4H_2O + Mn^{2+} E^\circ = 1.51V$

$$E^{\circ}$$
cell = 0.54 - 1.51 = -0.97V
E < 0 not spontaneous. Electrolytic cell!